

ORAL PRESENTATION

Open Access

# End growth results analysis related to Risser score, Cobb degrees, and curve types at the beginning of the treatment

S Donzelli<sup>1\*</sup>, F Zaina<sup>1</sup>, S Negrini<sup>2,3</sup>

From 9th International Conference on Conservative Management of Spinal Deformities - SOSORT 2012 Annual Meeting  
Milan, Italy. 10-12 May 2012

## Background

Scoliosis treatment is the science of prediction and estimation. The strategy is based on the magnitude, and pattern of the deformity, both related to age and Risser Score, to predict the potential progression. The efficacy of the conservative treatment of scoliosis is known, and analysis of factors potentially influencing the results is surely interesting.

## Aim

To assess the final results stratified according to curve magnitude, Risser score, curves type, gender and age, of a prospective set of patients treated in a centre fully dedicated to the conservative treatment of Adolescent Idiopathic Scoliosis AIS.

## Method

Study Design: Retrospective study. Population. 388 (31 males) patients respecting these inclusion criteria: AIS diagnosis, Risser test 0-3; all Cobb degrees; no prior treatment; who had reached the end of treatment since our institute database start in 2003. Methods: Clinical and radiographic (Cobb degrees) data at the beginning of treatment have been compared to end growth results. Treatments: All patients were treated respecting SOSORT standard of conservative treatment with observation, exercises, soft and rigid braces.

## Results

At the start of treatment worst curves corresponded to highest Risser test. With treatment, the percentage of

unchanged patients remained almost stable (40-50%), while progression was higher in Risser 0 than at other Risser stages (16.6% vs 7.4-9.9%); highest rates of improvement appeared at Risser 1 or 3 (45-47% vs 33-39%). Patients who begin the therapy at Risser 0 have a higher probability to end treatment under 30° Cobb. The probability of curve progression is highest at the lowest and highest initial sizes of curve (i.e. below 20° or over 40°). In this sample of treated patients, the rate of progressed curves was very low, with a high rate of stabilized and improved curves. The total number of patients who finished with curves over 30°C increases proportionally with a starting Risser Score. Considering curve's type, age and gender we didn't find differences in final results.

## Conclusion

The efficacy of conservative treatment has been demonstrated in some previous studies. Our data confirms this aspect, with a trend of all final results (82.6% below 30°C), telling us that therapy is the most important predictive factor. As expected, the lower the age and Risser at start the lower the curve magnitude, and the best the final results, confirming the importance of early AIS detection.

## Author details

<sup>1</sup>ISICO (Italian Scientific Spine Institute), Milan, Italy. <sup>2</sup>University of Brescia, Brescia, Italy. <sup>3</sup>IRCCS Don Gnocchi, Milan, Italy.

Published: 3 June 2013

## References

1. Coillard C, Vachon V, Circo AB, Beausejour M, Rivard CH: Effectiveness of the SpineCor brace based on the new standardized criteria proposed by the scoliosis research society for adolescent idiopathic scoliosis. *J Pediatr Orthop* 2007, **27**(4):375-379.

<sup>1</sup>ISICO (Italian Scientific Spine Institute), Milan, Italy  
Full list of author information is available at the end of the article

2. Lonstein JE, Carlson JM: **The prediction of curve progression in untreated idiopathic scoliosis during growth.** *J Bone Joint Surg Am* 1984, **66**(7):1061-1071.
3. Negrini S, Atanasio S, Zaina F, Romano M, Parzini S, Negrini A: **End-growth results of bracing and exercises for adolescent idiopathic scoliosis. Prospective worst-case analysis.** *Stud Health Technol Inform* 2008, **135**:395-408.

doi:10.1186/1748-7161-8-S1-O10

**Cite this article as:** Donzelli *et al.*: End growth results analysis related to Risser score, Cobb degrees, and curve types at the beginning of the treatment. *Scoliosis* 2013 **8**(Suppl 1):O10.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

