

GEOGRAPHIC, PERSONAL AND CLINICAL DETERMINANTS OF BRACE-WEARING TIME IN ADOLESCENTS WITH IDIOPATHIC SCOLIOSIS

Alessandra Negrini¹, Sabrina Donzelli², Giulia Rebagliati¹, Giulia Fregna³, Michele Romano¹, Fabio Zaina¹ and Stefano Negrini^{4,5}

1. ISICO (Italian Scientific Spine Institute), Milan, Italy
 2. IRCCS Istituto Auxologico Italiano, Italy
 3. Doctoral Program in Translational Neurosciences and Neurotechnologies, University of Ferrara, 44121 Ferrara, Italy.
 4. Department of Biomedical, Surgical and Dental Sciences, University of Milan“La Statale”, 20122 Milan, Italy
 5. IRCCS Galeazzi S.Ambrogio Hospital, 20157 Milan, Italy
- Purpose: Adherence to brace treatment is crucial for its effectiveness in AIS. The study aimed to verify the impact of geographic, personal and clinical variables on adherence to brace wear.

Methods: Design: retrospective cross-sectional study of consecutively collected participants at a tertiary referral institute. Inclusion criteria: AIS, one-year of follow-up after bracing, and use of a thermal sensor. The outcome was adherence and tested predictors were age, gender, ATR, TRACE, Cobb, curve type, Risser, and back pain at baseline, brace type and prescription time, income, living in the North, Centre, or South of Italy, city of residence size, distance from the sea, and height above sea level. The chi 2 test was used to investigate the association between the categories and adherence. The logistic regression model tested the predictors. STATA 17.0 MP was used.

Results: We included 1904 adolescents, age 13.8±1.6 (1597 females), worse scoliosis curve 35±11° Cobb. The prevalence of good compliers, with more than 75% of the prescribed dosage, was 90.8%. Factors associated with higher adherence were living in the North of Italy (p=0.003), prescription >20 hours per day (p=0.008), female gender (p=0.008), younger age and lower Risser sign (age 10 to 13: p=0.000; Risser 0-1: p=0.013). The regression model explained 4% of the variability.

Conclusion: Living in the North of Italy, > hours of brace prescription, female gender, younger age, and lower Risser sign increased adherence to brace wear. Nevertheless, there are other unknown factors that contribute to the explanation of adherence in this highly compliant population.